



Definition and diagnosis of fracture-related infection

Martin McNally¹
Geertje Govaert²
Maria Dudareva¹
Mario Morgenstern³
Willem-Jan Metsemakers⁴

- Fracture-related infection (FRI) is common and often diagnosed late.
- Accurate diagnosis is the beginning of effective treatment.
- Diagnosis can be difficult, particularly when there are no outward signs of infection.
- The new FRI definition, together with clear protocols for nuclear imaging, microbiological culture and histological analysis, should allow much better study design and a clearer understanding of infected fractures.
- In recent years, there has been a new focus on defining FRI and avoiding non-specific, poorly targeted treatment. Previous studies on FRI have often failed to define infection precisely and so are of limited value. This review highlights the essential principles of making the diagnosis and how clinical signs, serum tests, imaging, microbiology, molecular biology and histology all contribute to the diagnostic pathway.

Keywords: definition; diagnosis; fracture; fracture-related infection (FRI); histopathology; medical imaging; microbiology; serum inflammatory markers

Cite this article: *EFORT Open Rev* 2020;5:614-619.
DOI: 10.1302/2058-5241.5.190072

Introduction

Successful treatment of a fracture is aimed at promoting bone healing and good functional recovery with the avoidance of complications which may impair function or prolong treatment. When an infection occurs, this can be a devastating event, with the need for unexpected surgery, increased hospital stay and much higher health-care costs.^{1,2} The effects on the patient extend far beyond

the initial treatment of the infection or fracture. Patients may require support from healthcare providers and social services over a prolonged period.^{2,3} It is even more stressful for patients if the diagnosis of an infection is delayed or missed. Failure to begin treatment promptly may convert a simple early fracture infection into a chronic persistent infected non-union which may be very difficult to eradicate.^{4,5}

Considering the importance of diagnosing and treating an infected fracture, it is surprising that it is only very recently that an accepted definition of fracture-related infection (FRI) has been published. In 1996, Arens et al reported that in all studies they had reviewed, the term 'infection' was not defined.⁶ This situation remained unchanged for 22 years when, in 2018, Metsemakers et al found that only 2% of randomized trials of fracture fixation used any definition of infection when reporting this complication.⁷

By contrast to prosthetic joint infection, the lack of a definition or diagnostic strategy for infection has hampered the development of treatment protocols with comparable studies and outcomes. This deficit has prompted many surgeons to use diagnostic criteria developed for prosthetic joint infection. However, this is not appropriate, as the patient populations are different (elective arthroplasty versus traumatized patients) and pre-operative diagnostic tests from joint puncture or biopsy are often not feasible.

This review aims to summarize the significant advances which have been made in recent years towards establishing good diagnostic pathways with validated investigations. These have been developed by several groups around Europe and have been brought together through the FRI Consensus Group,⁸ the European Bone and Joint Infection Society (EBJIS), the AO Foundation, Pro-Implant and the Orthopaedic Trauma Association (OTA).




Fig. 1 The diagnostic algorithm for a suspected fracture-related infection (FRI), using the updated definition criteria of the FRI Consensus Group.²⁸

Note. NP/HPF, neutrophils per high-power field (x 400 magnification).

1. Any of these criteria will diagnose the presence of an infection independently. For instance, the presence of a draining sinus communicating with the fracture does not need microbiological culture for confirmation of infection.
2. Suggestive criteria should prompt the surgeon to consider further investigations which may lead to identifying possible confirmatory criteria.
3. A single positive culture from sonication fluid at >50 colony-forming units per ml, particularly from a virulent organism, such as *Staph. aureus*, is highly suggestive of infection and may be taken as confirmatory on an intention-to-treat basis.
4. >5NP/HPF is regarded as confirmatory of infection in cases more than 8 weeks after fracture.

Terminology and definition of infection

Many terms have been used to describe the onset of infection after fracture. The use of words like osteomyelitis, osteitis or deep surgical site infection can be confusing and do not give specific indication of the relationship with a fracture. The FRI Consensus Group published the initial definition criteria in 2018⁸ and the term 'fracture-related infection' was adopted to encompass all infections which occur in the presence of a fracture. This includes early infection around fracture implants, infected non-unions, haematogenous infections arising after fracture healing and infections in fractures with no internal fixation.

Review of the literature showed that there are some diagnostic tests which are highly specific for the presence of infection (confirmatory criteria). These included sinus tracks communicating with the fracture, microbiological culture of organisms and histological features of infection. Conversely, there were features which suggested the presence of infection but could not be diagnostic alone

(suggestive criteria). These included some clinical signs, blood biomarkers and imaging tests. The FRI definition has recently been updated to include new data and is summarized in Fig. 1.

Blood biomarkers

Common serum inflammatory markers such as leukocyte count, C-reactive protein and erythrocyte sedimentation rate have been evaluated in the diagnosis of FRI. All are non-specific and can be raised after trauma without infection and in many other inflammatory conditions. Also, they can be normal in many chronic or late infections. C-reactive protein (CRP) levels rise after injury to a maximum at day 2 and then reduce to normal over 1–2 weeks.⁹ Systematic review of the literature reveals that CRP is perhaps the most useful marker, but with only moderate sensitivity and specificity.¹⁰ In recent studies of infected fractures and non-unions, blood markers had very limited predictive value.^{11,12} They cannot be used to confirm or exclude an infection.

Imaging for FRI

Imaging is inextricably linked to fracture surgery. When a post-operative FRI is considered, there are three main indications to (re)image the affected limb:

1. To evaluate fracture consolidation and implant stability.
2. To determine if an infection is present.
3. To assess the extent of that infection with specific anatomical details (such as sinuses, sequestra and cloacae) for surgical planning.

The most commonly used imaging modalities are plain X-ray, computed tomography (CT), magnetic resonance imaging (MRI) and the nuclear imaging techniques of white blood cell (WBC) scintigraphy and fluorodeoxyglucose positron emission tomography (^{18}F FDG-PET). Three-phase bone scintigraphy (BS) is highly sensitive for detecting FRI (89–100%) but unfortunately has a very low specificity (0–10%).¹³ BS is therefore almost obsolete for this indication.

In established FRI, for example in the presence of a draining sinus,⁸ imaging should focus on determining fracture consolidation and implant stability. For this purpose, plain X-ray, possibly combined with a CT scan, is usually sufficient. If the presence of an infection or the extent of the infection is the issue, more advanced imaging techniques such as MRI or nuclear imaging are required. MRI has many advantages: it is familiar, relatively cheap and widely available. Also it can distinguish anatomical details in the bone and adjacent soft tissues and has excellent sensitivity for detecting FRI (82–100%).¹³ With artifact reduction techniques the interference of metal implants can be reduced to a minimum.¹⁴ The downside of MRI is that, probably due to its inability to differentiate between sterile inflammation, normal bone healing and infected tissue, its specificity is reduced (43–60%).¹³

Nuclear imaging is far more accurate in cases where it is important to distinguish infected from non-infected tissues. It has been shown that both WBC scintigraphy and ^{18}F FDG-PET are highly accurate for diagnosing FRI shortly after surgery.^{15,16} More recently, the use of hybrid camera systems which combine nuclear imaging with CT (single photon emission computed tomography (SPECT) (SPECT/CT, PET/CT or PET/MRI) has led to increased accuracy and better anatomical details.^{13,17–19} Large retrospective studies show a sensitivity and specificity of WBC scintigraphy with SPECT/CT for diagnosing FRI, of 79–100% and 97% respectively.^{15,20} For ^{18}F FDG-PET, sensitivity is 88–89% and specificity is 76–80%.^{16,21} This means that WBC scintigraphy + SPECT/CT is slightly more accurate than ^{18}F FDG-PET/CT, but one has to bear in mind that all these studies were

hampered by their retrospective study designs. Secondly, although less accurate, ^{18}F FDG-PET has major advantages over WBC scintigraphy in terms of lower complexity of the labelling procedure, the requirement for just one scan, rather than early and late phase scans (over 20 hours), and its higher spatial resolution (3–4 mm vs. 8 mm).¹⁹

For the diagnosis of FRI, every imaging modality has its advantages and disadvantages and most studies have methodical limitations. A prospective clinical trial has recently been launched that will hopefully provide more clarity about the optimal imaging strategy for FRI in terms of diagnostic accuracy, cost-effectiveness and how to solve logistic challenges.²² While awaiting these results, the imaging modality of choice therefore mostly depends on the clinical question to be answered and local availability and experience with the imaging technique of choice.

Microbiological diagnosis of FRI

Published evidence relating to the microbiological diagnosis of FRI should be considered with the caveat that the consensus definition of FRI is recent, and that many previous studies excluded patients on the basis of the duration, site, and other clinical features of infection.

The aims of microbiological diagnosis of FRI are two-fold:

1. To confirm fracture-related infection when it is suspected.
2. To identify the infecting pathogen(s) and assess antimicrobial susceptibility patterns, and to select a targeted and tolerable, preferably oral, antimicrobial treatment for the patient.

Confirmation of infection

Accurate microbiological diagnosis requires the analysis of representative, uncontaminated samples of tissue or fluid from the fracture. Pre-operative diagnosis from superficial swabs, biopsies or aspirations is usually unhelpful as there is often poor concordance between biopsies and intra-operative sampling. A high proportion of FRIs are caused by bacteria that may be considered skin commensals. Detection of organisms such as coagulase-negative Staphylococci, Corynebacteria, and *Cutibacterium acnes* on a single biopsy specimen is thus very difficult to interpret. Also, up to one third of patients with FRI have polymicrobial infection, which is often missed in pre-operative biopsies.^{23,24} If antimicrobial therapy is based on limited pre-operative cultures alone, it has been shown that poor outcomes will be achieved.²⁵

Reliable microbiological information is obtained by culture of at least five uncontaminated samples, collected during surgery. Samples should be collected in a structured process, with separate instruments for each sample

and without touching the patient's skin with the sample or instrument.^{24,26,27} A study of the implementation of a structured sampling protocol following the recommendations for the diagnosis of FRI, found a greater proportion of polymicrobial infections diagnosed after the implementation, compared with unstructured microbiological sampling prior to the implementation, despite no increase in the proportion of positive samples or suspected culture contamination.²⁴ Specimens should be transferred rapidly to the laboratory and handled in a standardized protocol.²⁸ Five operative specimens can be processed under aerobic and anaerobic conditions with selective and non-selective media for less than 55 Euros per patient.²⁹ Automated liquid culture may be even more cost-effective.^{30,31}

Infections around implants or dead bone will frequently occur in the presence of biofilm. Methods which disrupt biofilm may enhance the isolation of organisms and improve diagnostic yield.²⁸ Sonication is a useful adjunct in the diagnosis of FRI where implants or cortical bone fragments are removed during operative debridement. The sensitivity of sonication alone in a series of 158 fractures, assessed against a clinical diagnosis of infection (sinus, purulence or histopathology findings consistent with infection) was 68%.³² Prior studies reported sonication fluid sensitivity for FRI at between 65% and 95%, with specificity between 50% and 97%.³³ Sonication should be used together with deep tissue sampling for optimum diagnostic accuracy.^{28,32}

Microbiological confirmation of FRI is obtained when phenotypically indistinguishable organisms are cultured from at least two separate deep tissue/implant specimens (including sonication fluid), collected and processed as above. A single positive culture is not confirmatory but is suggestive of infection and should be confirmed with other tests.^{8,28}

Identification of pathogens and antimicrobial susceptibilities

FRI is caused by a very wide variety of bacteria and fungi. Additionally, polymicrobial infection is common and often includes Gram-negative organisms.^{23,34} Clinical characteristics (open fracture, presence of metalwork, draining sinus, compromised host) do not discriminate sufficiently to anticipate who may be affected with polymicrobial or non-staphylococcal FRI.²³ The microbiological diagnosis of FRI must involve methods which determine antimicrobial susceptibility in individual microbial species, as antimicrobial sensitivity cannot be reliably predicted. Up to one third of classifiable organisms causing FRI may be multi-drug-resistant (MDR), even in countries with a low incidence of MDR infection.²³ The presence of these resistant bacteria adversely affects outcome.³⁵ Therefore, there is no place for empirical broad-spectrum therapy, particularly aimed at Gram-positive bacteria.

Antibiotic side-effects are frequent in the context of FRI treatment for at least six weeks, with around one in six patients affected, so the careful choice of antimicrobials for FRI must be justified by microbiological diagnosis.³⁶ The use of non-targeted, broad-spectrum agents is associated with subsequent multi-resistant infection for individual patients^{37,38} and cannot be justified. Intravenous empirical therapy should only be used immediately after surgery, while microbiological diagnosis is awaited. Thus, antimicrobial susceptibility assessment must form an integral part of the microbiological diagnosis of FRI.

Molecular methods have now been investigated in the diagnosis of FRI. Whole genome sequencing³⁹ and multiplex polymerase chain reaction methods (PCR)⁴⁰ have shown initial encouraging results. PCR-based identification panels and 16s-PCR have low sensitivity for the detection of all infecting organisms and may not be able to detect fungal or unusual bacterial infection. An evaluation of sonication fluid PCR for the microbiological diagnosis of FRI found that *Candida spp.* and *Enterococcus spp.*, pathogens with particularly poor treatment outcomes in the context of orthopaedic infection, were especially likely to be missed.⁴⁰ Currently, molecular techniques cannot give useful information on antimicrobial susceptibility.

Histopathology

The presence of visible microorganisms in deep tissue, as confirmed by histopathological examination using specific staining techniques for bacteria (e.g. Gram stain, Ziehl-Neelsen stain for tuberculosis or Grocott methenamine silver stain for fungi), is regarded as a confirmatory sign of FRI.⁸ In prosthetic joint infection (PJI), the presence of ≥ 5 polymorphonuclear neutrophils per high-power field (PMN/HPF) in five high-power fields observed in histological sections, at x 400 magnification, is considered to be an important intra-operative criterion for PJI.⁴¹ In contrast to the definition for PJI, the FRI Consensus Group did not include the presence of an acute inflammatory cell infiltrate on histopathological examination (i.e. PMN count) in the first version of the FRI definition.⁸ The reason for this was the lack of clear scientific evidence and, more specifically, agreement on a cut-off value above which FRI can be reliably diagnosed.^{8,33}

However, a large study on the value of quantitative histopathology for the diagnosis of chronic/late-onset FRI (at least two months after fracture) has now been published.⁴² Morgenstern et al. proposed a novel bimodal approach to confirm or exclude infection in unhealed fractures. The complete absence of PMNs had a very high correlation with aseptic non-union (specificity 98%, positive predictive value (PPV) 98%). Conversely, the presence of >5 PMN/HPF only occurred with infection (specificity 100%; PPV 100%). The combination of clinical signs, ≥ 2

microbiological cultures and bimodal histopathological analysis (absent NPs vs. >5 PMN/HPF) improved diagnostic accuracy in up to 96.8% of cases. Due to increasing evidence, the presence of >5 PMN/HPF in fractures more than 8 weeks after injury (e.g. fracture non-union) was added by the expert panel in the second consensus meeting as the fifth confirmatory sign of FRI.²⁸ (Fig. 1).

Conclusions

Interest in the definition and diagnosis of FRI is a recent development. There are now good studies which give clear definition criteria and some evidence around the effectiveness of diagnostic methods. Pre-operative diagnosis with serum markers, superficial swabs or percutaneous biopsies is not accurate. Diagnosis depends on the interpretation of deep tissue specimens for microbiological culture and histological analysis. The improvements in imaging, particularly with nuclear medicine combined with localizing scans (¹⁸F-DG-PET/CT or SPECT/CT), offer new possibilities for the diagnosis of infection and for planning of surgical procedures.

There is now no place for empirical treatment of suspected infection, in the hope of a miraculous cure. Careful attention to establishing the diagnosis allows better surgical planning and pathogen-specific antimicrobial therapy, designed to improve outcomes in FRI.

AUTHOR INFORMATION

¹The Oxford Bone Infection Unit, Nuffield Orthopaedic Centre, Oxford University Hospitals, Oxford, UK.

²Department of Trauma Surgery, University of Utrecht, University Medical Center Utrecht, Utrecht, The Netherlands.

³Department of Orthopaedic and Trauma Surgery, University Hospital Basel, Switzerland.

⁴Department of Trauma Surgery, University Hospitals Leuven, Leuven, Belgium.

Correspondence should be sent to: Martin McNally, Oxford Bone Infection Unit, Nuffield Orthopaedic Centre, Windmill Road, Oxford, OX3 7HE, UK.
Email: martin.mcnally@ouh.nhs.uk

ICMJE CONFLICT OF INTEREST STATEMENT

MM reports textbook royalties from Oxford University Press; expenses paid for travel and accommodation to attend symposia or expert panels from Bonesupport AB, outside the submitted work.

GG reports being a board member of the Dutch Trauma Society, and a research grant for prospective imaging trial (the IFI trial) from DePuy-Synthes, outside the submitted work.

The other authors declare no conflict of interest relevant to this work.

FUNDING STATEMENT

No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article.

LICENCE

© 2020 The author(s)

This article is distributed under the terms of the Creative Commons Attribution-Non Commercial 4.0 International (CC BY-NC 4.0) licence (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed.

REFERENCES

- Olesen UK, Pedersen NJ, Eckardt H, et al.** The cost of infection in severe open tibial fractures treated with a free flap. *Int Orthop* 2017;41:1049–1055.
- Parker B, Petrou S, Masters JPM, Achana F, Costa ML.** Economic outcomes associated with deep surgical site infection in patients with an open fracture of the lower limb. *Bone Joint J* 2018;100-B:1506–1510.
- Tutton E, Achten J, Lamb SE, Willett K, Costa ML.** A qualitative study of patient experience of an open fracture of the lower limb during acute care. *Bone Joint J* 2018;100-B:522–526.
- McNally M, Ferguson J, Kugan R, Stubbs D.** Ilizarov treatment protocols in the management of infected nonunion of the tibia. *J Orthop Trauma* 2017;31:S47–S54.
- Bose D, Kugan R, Stubbs D, McNally M.** Management of infected nonunion of the long bones by a multidisciplinary team. *Bone Joint J* 2015;97-B:814–817.
- Arens S, Hansis M, Schlegel U, et al.** Infection after open reduction and internal fixation with dynamic compression plates: clinical and experimental data. *Injury* 1996;27:SC27–SC33.
- Metsemakers WJ, Kortram K, Morgenstern M, et al.** Definition of infection after fracture fixation: a systematic review of randomized controlled trials to evaluate current practice. *Injury* 2018;49:497–504.
- Metsemakers WJ, Morgenstern M, McNally MA, et al.** Fracture-related infection: a consensus on definition from an international expert group. *Injury* 2018;49:505–510.
- Neumaier M, Scherer MA.** C-reactive protein levels for early detection of postoperative infection after fracture surgery in 787 patients. *Acta Orthop* 2008;79:428–432.
- van den Kieboom J, Bosch P, Plate JDJ, et al.** Diagnostic accuracy of serum inflammatory markers in late fracture-related infection: a systematic review and meta-analysis. *Bone Joint J* 2018;100-B:1542–1550.
- Bosch P, van den Kieboom J, Plate JDJ, et al.** Limited predictive value of serum inflammatory markers for diagnosing fracture-related infections: results of a large retrospective multicenter cohort study. *J Bone Jt Infect* 2018;3:130–137.
- Sigmund IK, Morgenstern M, Dudareva M, Athanasou N, McNally MA.** Limited diagnostic value of serum inflammatory biomarkers in the diagnosis of fracture-related infections. *Bone Joint J* in press.
- Govaert GA, Ijpma FFA, McNally M, McNally E, Reininga IH, Glaudemans AW.** Accuracy of diagnostic imaging modalities for peripheral post-traumatic osteomyelitis: a systematic review of the recent literature. *Eur J Nucl Med Mol Imaging* 2017;44:1393–1407.
- Gupta A, Subhas N, Primak AN, Nittka M, Liu K.** Metal artifact reduction: standard and advanced magnetic resonance and computed tomography techniques. *Radiol Clin North Am* 2015;53:531–547.
- Govaert GAM, Bosch P, Ijpma FFA, et al.** High diagnostic accuracy of white blood cell scintigraphy for fracture related infections: results of a large retrospective single-center study. *Injury* 2018;49:1085–1090.
- Lemans JVC, Hobbelink MGG, Ijpma FFA, et al.** The diagnostic accuracy of ¹⁸F-FDG PET/CT in diagnosing fracture-related infections. *Eur J Nucl Med Mol Imaging* 2019;46:999–1008.

- 17. Glaudemans AWJM, Bosch P, Slart RHJA, IJpma FFA, Govaert GAM.** Diagnosing fracture-related infections: can we optimize our nuclear imaging techniques? *Eur J Nucl Med Mol Imaging* 2019;46:1583–1587.
- 18. Glaudemans AW, Prandini N, Di Girolamo M, et al.** Hybrid imaging of musculoskeletal infections. *Q J Nucl Med Mol Imaging* 2018;62:3–13.
- 19. Govaert GAM, Glaudemans AWJM.** Nuclear medicine imaging of posttraumatic osteomyelitis. *Eur J Trauma Emerg Surg* 2016;42:397–410.
- 20. Glaudemans AW, de Vries EF, Vermeulen LE, Slart RH, Dierckx RA, Signore A.** A large retrospective single-centre study to define the best image acquisition protocols and interpretation criteria for white blood cell scintigraphy with ^{99m}Tc-HMPAO-labelled leucocytes in musculoskeletal infections. *Eur J Nucl Med Mol Imaging* 2013;40:1760–1769.
- 21. Wenter V, Müller JP, Albert NL, et al.** The diagnostic value of [(18)F]FDG PET for the detection of chronic osteomyelitis and implant-associated infection. *Eur J Nucl Med Mol Imaging* 2016;43:749–761.
- 22. Govaert G, Hobbelink M, Reininga I, et al.** The accuracy of diagnostic imaging techniques in patients with a suspected fracture-related infection (IFI) trial: study protocol for a prospective multicenter cohort study. *BMJ Open* 2019;9:e027772.
- 23. Dudareva M, Hotchen AJ, Ferguson J, et al.** The microbiology of chronic osteomyelitis: changes over ten years. *J Infect* 2019;79:189–198.
- 24. Hellebrekers P, Rentenaar RJ, McNally MA, et al.** Getting it right first time: the importance of a structured tissue sampling protocol for diagnosing fracture-related infections. *Injury* 2019;50:1649–1655.
- 25. Taormina DP, Shulman BS, Lee JH, Karia RJ, Marciano AI, Egol KA.** The outcome of patients with cultured pathogens at time of nonunion surgery. *Acta Orthop Belg* 2018;84:1–10.
- 26. Dudareva M, Barrett L, Morgenstern M, et al.** An evidence base for tissue sampling and culture interpretation in fracture-related infection. *Bone Joint J Orthopaedic Proceedings* 2018;100-B.
- 27. Sigmund IK, McNally MA.** Diagnosis of bone and joint infections. *Orthop Trauma* 2019;33:144–152.
- 28. Govaert GAM, Kuhl R, Atkins BL, et al.** Diagnosing fracture-related infections: current concepts and recommendations (FRI Consensus Group). *J Orthop Trauma* 2020;34:8–17.
- 29. Bémer P, Léger J, Tandé D, et al; Centre de Référence des Infections Ostéo-articulaires du Grand Ouest (CRIOGO) Study Team.** How many samples and how many culture media to diagnose a prosthetic joint infection: a clinical and microbiological prospective multicenter study. *J Clin Microbiol* 2016;54:385–391.
- 30. Minassian AM, Newnham R, Kalimeris E, Bejon P, Atkins BL, Bowler ICJW.** Use of an automated blood culture system (BD BACTEC™) for diagnosis of prosthetic joint infections: easy and fast. *BMC Infect Dis* 2014;14:233.
- 31. Peel TN, Sedarski JA, Dylla BL, et al.** Laboratory workflow analysis of culture of periprosthetic tissues in blood culture bottles. *J Clin Microbiol* 2017;55:2817–2826.
- 32. Dudareva M, Barrett L, Figtree M, et al.** Sonication versus tissue sampling for diagnosis of prosthetic joint and other orthopaedic device-related infections. *J Clin Microbiol* 2018;56:e00688–18.
- 33. Onsea J, Depypere M, Govaert G, et al.** Accuracy of tissue and sonication fluid sampling for the diagnosis of fracture-related infection: a systematic review and critical appraisal. *J Bone Jt Infect* 2018;3:173–181.
- 34. Otchwemah R, Moczko T, Marche B, Mattner F, Probst C, Tjardes T.** High prevalence of bacteria in clinically aseptic non-unions of the tibia and the femur in tissue biopsies. *Eur J Trauma Emerg Surg* 2018. [Epub ahead of print].
- 35. Hotchen A, Dudareva M, Ferguson J, Rombach I, Scarborough M, McNally M.** Does the BACH Classification of long bone osteomyelitis correlate with patient reported outcome measures following surgery? *Bone Joint Res* 2019;8:459–468.
- 36. Conterno LO, Turchi MD.** Antibiotics for treating chronic osteomyelitis in adults. *Cochrane Database Syst Rev* 2013;9:CD004439.
- 37. Defez C, Fabbro-Peray P, Bouziges N, et al.** Risk factors for multidrug-resistant *Pseudomonas aeruginosa* nosocomial infection. *J Hosp Infect* 2004;57:209–216.
- 38. Doi Y, Park YS, Rivera JI, et al.** Community-associated extended-spectrum β -lactamase-producing *Escherichia coli* infection in the United States. *Clin Infect Dis* 2013;56:641–648.
- 39. Street TL, Sanderson ND, Atkins BL, et al.** Molecular diagnosis of orthopaedic-device-related infection directly from sonication fluid by metagenomic sequencing. *J Clin Microbiol* 2017;55:2334–2347.
- 40. Renz N, Cabric S, Morgenstern C, Schuetz MA, Trampuz A.** Value of PCR in sonication fluid for the diagnosis of orthopedic hardware-associated infections: has the molecular era arrived? *Injury* 2018;49:806–811.
- 41. Parvizi J, Tan TL, Goswami K, et al.** The 2018 definition of periprosthetic hip and knee infection: an evidence-based and validated criteria. *J Arthroplasty* 2018;33:1309–14 e2.
- 42. Morgenstern M, Athanasou NA, Ferguson JY, Metsemakers WJ, Atkins BL, McNally MA.** The value of quantitative histology in the diagnosis of fracture-related infection. *Bone Joint J* 2018;100-B:966–972.